PRIDE MOBILITY®

Educational Clinical Evaluation and Documentation Resource for Power Lift Recliners

Important Notice to Health Care Professionals and Providers Regarding this Resource's Educational Purpose and the Independence of Clinical Judgment

This resource is provided solely for general educational and informational purposes. It does not recommend treatment for any individual patient, prescribe or direct the use of any particular device or modality, establish a standard of care, dictate documentation, or in any way steer, control, influence, or substitute for the independent clinical judgment of any physician, occupational therapist, physical therapist, provider, or other licensed health care professional. Nothing in this resource is intended to direct, steer, or improperly influence any clinician's independent patient assessment, diagnosis, treatment recommendation, coverage determination, or medical-necessity determination. Each clinician remains solely responsible for independently evaluating each patient, for determining whether and to what extent any assistive device is clinically appropriate based on his or her own professional judgment, for documenting only findings that accurately reflect the clinician's own examination, and for complying with all applicable payer requirements and professional standards.

1. Purpose

This clinical evaluation and documentation resource is intended to provide educational information that may assist health care professionals in considering whether a power lift recliner may be relevant to, or may support, a patient's functional limitations, subject always to the health care professional's independent, individualized evaluation and assessment of each patient. [1; 3; 9; 13; 14; 15]

2. Illustrative Clinical Documentation Examples (Examples Only)

The following examples are provided for illustrative and educational purposes only. They are not templates, model language, required language, or suggested chart entries, and they must never be copied verbatim into a patient's medical record. All documentation must reflect the clinician's own independent evaluation, professional judgment, and patient-specific findings, and nothing in these examples is intended to recommend, direct, steer, or otherwise influence any clinician's assessment, diagnosis, treatment recommendation, or medical-necessity determination.

Example: *Following an individualized professional evaluation, a clinician may determine that a power lift recliner/electric positioning chair could support or assist a patient who has been found unable to safely and independently rise from a standard household armchair in connection with a diagnosed medical condition and corresponding symptomatology. In such circumstances, the lift chair's powered seat-lift mechanism may assist with a smooth, controlled rise from seated to near-standing, which may help reduce caregiver lifting assistance and support reduced transfer-related fall risk during sit-to-stand and stand-to-sit transitions, subject always to the clinician's individualized evaluation. Once standing, the patient is able to ambulate/transfer using an assistive device to complete MRADLs such as toileting, dressing, hygiene, meals, and access to the home environment. [1; 2; 3; 7; 9; 10; 11; 12; 13; 14]*

PRIDE MOBILITY®

Power Lift Recliner Features That May Be Relevant to Certain Forms of Mobility Impairment

Product Feature	Potential clinical relevance (subject to independent clinical evaluation)
Powered seat-lift mechanism/ stand assist	May support sit-to-stand and stand-to-sit by tilting/elevating the seating surface for patients with reduced lower-extremity strength, pain, impaired balance, poor endurance, or unsafe transfer mechanics. [1; 2; 7; 8; 9; 10; 11; 12; 14]
Remote-controlled operation	May support independent operation when the patient has the cognition, vision, reach, and hand function needed to use the hand control safely; supervision or controller-access plan must be documented when identified. [4]
True-infinite positioning/ independent backrest and footrest movement	May allow individualized positioning for postural support, seated tolerance, lower-extremity elevation, rest breaks, and recovery positioning when clinically justified. [18]
Power headrest and power lumbar	May support head, neck, and trunk alignment and seated tolerance when consistent with the clinician's documented postural findings, subject to independent clinical evaluation. [18]

PRIDE MOBILITY®

Clinical Circumstances in Which a Power Lift Recliner May Be Relevant

A. Unsafe or Dependent Sit-to-Stand from Standard Seating

- A power lift recliner may be relevant where the clinician, following an independent and individualized evaluation, determines that a patient cannot rise from a standard armchair without physical assistance, excessive compensatory movement, multiple attempts, or unsafe use of furniture. [1; 2; 8; 9; 14]
- Clinical consideration of a power lift recliner may be appropriate where the clinician identifies and documents a history of chair height issues, increased need for mobility assistance, pain, weakness, range of motion limits, balance deficits, endurance limits, fall history, where the patient can ambulate after standing. [1; 2; 7; 8; 9]

> ***Example:*** *Following evaluation, the clinician determines that the patient is unable to rise from a standard 18-inch-high armchair without moderate physical assistance due to bilateral knee pain, decreased quadriceps strength, and impaired anterior weight shift. Use of the powered seat-lift mechanism in a power lift recliner allows controlled elevation to near-standing, reducing caregiver lifting and enabling the patient to stand and ambulate with a rolling walker to access toileting and hygiene tasks. [1; 2; 7; 9; 14; 15]*

B. Controlled Stand-to-Sit Due to Poor Eccentric Control

- A power lift recliner may be relevant where the clinician, following an independent and individualized evaluation, determines that a patient drops into chair seating, loses balance when turning to sit, or cannot control descent due to lower-extremity weakness, pain, Parkinsonian movement patterns, neuropathy, stroke-related weakness, or fatigue. [2; 3; 9; 10; 11; 13]
- Clinical consideration of a power lift recliner may be appropriate where the clinician identifies and documents history of unsafe descent pattern, posterior fall risk, need for caregiver stabilization, and improvement with powered controlled seating. [2; 3; 9; 10; 13]

> ***Example:*** *Following evaluation, the clinician determines that the patient demonstrates poor eccentric control during stand-to-sit and falls backward into standard seating, creating risk for injury. A clinician may conclude that a power lift recliner provides controlled lowering from standing to seated position, supporting safer transfers and reducing need for caregiver stabilization. [2; 3; 9; 13]*

C. Severe Hip or Knee Arthritis Limiting Transfers

- A power lift recliner may be relevant where the clinician, following an independent and individualized evaluation, determines that a patient has functional transfer limitations associated with hip or knee osteoarthritis, rheumatoid arthritis, post-traumatic arthritis, or joint deformity that prevents standing from household armchairs. [6; 7; 8; 14]
- Clinical consideration of a power lift recliner may be appropriate where the clinician identifies and documents a history of severe hip/knee arthritis, inability to stand from any standard armchair in the home, failed conservative strategies, device as part of the treatment plan, and ability to ambulate once standing. [6; 8; 9]

Example: *Following evaluation, the clinician determines that due to severe bilateral knee osteoarthritis, the patient is unable to stand from any standard household armchair despite armrests and use of a walker. Once assisted to standing, the patient ambulates household distances with a rolling walker. A power lift chair may be medically appropriate and necessary to support safe chair-to-stand transitions needed for toileting, meals, and hygiene. [6; 8; 7; 14]*

D. Severe Neuromuscular Disease Affecting Transfer Independence

- A power lift recliner may be relevant where the clinician, following an independent and individualized evaluation, determines that a patient with Parkinson disease, muscular dystrophy, ALS, MS, post-polio syndrome, SCI, stroke-related weakness, or other neuromuscular impairment experiences functional transfer limitations as a result of the diagnosed condition, subject always to the clinician's determination of medical necessity. [5; 10; 11; 12; 14; 15]
- Clinical consideration of a power lift recliner may be appropriate where the clinician identifies and documents a history of lower-extremity weakness, rigidity, spasticity, bradykinesia, fatigue, poor trunk control, impaired balance, or similar significant mobility impairments, and based on their evaluation, concludes that standard seating adaptations are insufficient to promote safe sitting and standing. [2; 5; 8; 10; 11; 12; 14; 15]

Example: *Clinician's evaluation of the patient indicates that patient suffers from severe neuromuscular impairment resulting in reduced lower-extremity power, impaired trunk control, and inability to complete sit-to-stand from standard seating without caregiver lifting. The clinician may determine that a power lift recliner would provide a controlled elevation surface to compensate for impaired transfer biomechanics and thus would support safer sitting and standing and engagement with other household MRADLs. [5; 8; 10; 11; 12; 14; 15]*

E. Transfer-Related Fall-Risk Reduction

- A power lift recliner may be relevant where the clinician, following an independent and individualized evaluation, determines that a patient experiences falls or near-falls during chair transfers, and not only during walking. [3; 9; 13]
- Clinician identifies and documents a history of a number of falls and associated timing, notes a correlation to chair transfer phase where falls have occurred, notes environmental barriers, and may determine that a power lift chair may be appropriate to help alleviate some fall-related safety challenges for the patient. [3; 9; 13]

Example: *The clinician's evaluation of the patient suggests that the patient has experienced repeated near-falls when attempting to stand from household seating. The clinician may conclude that a powered lift mechanism would reduce the force and balance demands for the patient that are required to transition from sitting to standing, supporting safer initiation of gait that may be accomplished with an appropriate walker when used as evaluated and trained. [3; 7; 9; 13]*

F. Reduced Caregiver Physical Assistance and Injury Risk

- A power lift recliner may be appropriate where a clinician's evaluation concludes that a caregiver must pull, lift, boost, or physically transfer the patient from a chair. [1]
- Clinician identifies and documents a history of current caregiver assist at a high level and frequency reflecting some significant level of diminished independence for the patient, and where the caregiver has limitations. [1; 9]

  Example: *Clinician's evaluation indicates that the patient currently requires caregiver lifting assistance for every chair transfer multiple times per day. Clinician may determine that a powered lift chair would assist patient transfers to standing with supervision only, reducing caregiver physical lifting and improving access to toileting, meals, and mobility within the home. [1; 9]*

- A clinician determines after evaluation that a patient recovering from hip/knee surgery, spinal surgery, prolonged hospitalization, deconditioning, or other acute condition cannot safely rise from standard seating. [6; 14; 16; 17]
- Clinician identifies and documents a history of surgical precautions, pain, ROM limits, weakness, endurance deficits, expected duration of need if temporary, and relationship to home discharge safety, and determines that a power lift recliner may be an appropriate device to help address the functional limitations of the patient's diminished mobility. [6; 9; 14; 16; 17]

  Example: *Clinician's evaluation concludes, following hospitalization and significant deconditioning, that the patient is unable to rise from household seating without moderate assistance and further evidences unsafe compensatory movement. Clinician may determine that a power lift recliner supports safe home function by enabling controlled sit-to-stand practice and reducing caregiver lifting during the rehabilitation period. [6; 14; 15; 16; 17]*

Deficit Identification: When identified and properly documented by a clinician, the following patient evaluation-based findings may be relevant to a clinician's independent consideration of whether a power lift recliner is appropriate, subject always to individualized professional evaluation. Findings must always be based upon a clinician's independent evaluation of the patient. The following are not intended to prescribe treatment and should not be copied verbatim.

PRIDE MOBILITY®

Condition Based Evaluation: The following examples of conditions illustrate certain types of patient-specific findings that may correlate to a clinician's post evaluation determination that a power lift chair may be appropriate to assist in addressing certain mobility impairments. The references below to clinical conditions are illustrative only and are not intended to influence or interfere with a clinician's independent evaluation and conclusions, nor are they intended to prescribe a specific form of treatment or documentation, and should not be copied verbatim.

Condition-Based Examples

Condition/ presentation	Example clinical wording
Severe knee OA	Client has severe bilateral knee osteoarthritis with pain 8/10 during sit-to-stand, reduced knee extension strength, and inability to rise from standard seating without physical assistance. A power lift recliner may support reduced joint loading during transfer initiation and may assist with a safer transition to walker-assisted ambulation, subject to the clinician's independent evaluation. [6; 7; 8; 14]
Parkinson disease	Client demonstrates bradykinesia, freezing during transfer initiation, and impaired anterior weight shift. A power lift recliner may provide consistent, controlled elevation that can assist with sit-to-stand initiation and support reduced transfer-related fall risk during functional transfers, subject to the clinician's independent evaluation. [2; 3; 11; 12; 13]
Muscular dystrophy/ progressive neuromuscular weakness	Client has progressive proximal weakness and cannot generate sufficient hip/knee extension force from standard chair height. Power lift assistance may be relevant to preserving transfer independence and may assist with reducing caregiver lifting during daily MRADLs, subject to the clinician's independent evaluation. [5; 8; 9; 14; 15]
Stroke with hemiparesis	Client has hemiparesis, impaired balance, and unsafe push-off from standard seating. Powered lift and controlled descent may reduce transfer instability and may support safer access to standing mobility with a prescribed assistive device, subject to the clinician's independent evaluation. [2; 3; 10; 13]
Post-hospital deconditioning	Client presents with decreased endurance, generalized weakness, and inability to complete repeated sit-to-stand transfers from household seating without assistance. A lift recliner may support safe home function during recovery and may reduce caregiver physical assistance, subject to the clinician's independent evaluation. [6; 14; 15; 16; 17]
Control and driving ability	Doorway and turning access, surfaces, route length, charging, storage, and transport feasibility. [CER Stage 3; 3]
Environment	Actual performance with walking aids, seating/rest strategies, manual mobility, or other powered mobility as relevant. Explain why the selected option fits the findings. [CER Stage 3]
Alternatives considered	Tasks completed, assistance needed, errors, comfort, fatigue, and remaining limitations with the selected scooter. [CER Stage 3]
Trial results	Training provided, charging and maintenance arrangements, support available, and reassessment needs. [3]

PRIDE MOBILITY®

Source Links

- Pride Mobility - Lift Chair Recliners
- Pride Mobility - Lifting Recliners
- Pride Mobility - VivaLift! Ultra PLR-4955 product page
- Pride Mobility - VivaLift! Ultra overview
- Pride Mobility - Lift Chair Medicare Consumer Guide PDF
- Quantum Rehab - Medical Necessity for iLevel Power Adjustable Seat Height PDF
- CMS - Seat Lift Mechanisms LCD L33801
- CMS - Seat Lift NCD 280.4

Article References

[1] Edlich et al., "Revolutionary Advances in Adaptive Seating Systems for the Elderly and Persons with Disabilities That Assist Sit-to-Stand Transfers," National Center for Biotechnology Information, U.S. National Library of Medicine, 2003, https://www.ncbi.nlm.nih.gov/pubmed/?term=recline+lift+chair.

[2] Millington, Pamela, et al., "Biomechanical Analysis of the Sit-to-Stand Motion in Elderly Persons," Archives of Physical Medicine and Rehabilitation, July 1992, https://www.archives-pmr.org/article/0003-9993(92)90124-F/fulltext; Riley, P. O., et al., "Mechanics of a Constrained Chair-Rise," Journal of Biomechanics, 1991, https://www.ncbi.nlm.nih.gov/pubmed/2026635.

[3] Connell, Bettye R., and Steven L. Wolf, "Environmental and Behavioral Circumstances Associated with Falls at Home Among Healthy Elderly Individuals," Archives of Physical Medicine and Rehabilitation, Feb. 1997, https://www.archives-pmr.org/article/S0003-9993(97)90261-6/fulltext; Josama, Pascale, et al., "Poster 133: Cause of Injury and Mortality after Traumatic Brain Injury in an Older Population," Archives of Physical Medicine and Rehabilitation, Oct. 2011, https://doi.org/10.1016/j.apmr.2011.07.162.

[4] Geriatric and Long-Term Care Review Committee, "GLTCRC-2021-7A, 7B and 7C," https://www.coto.org/docs/default-source/default-document-library/appendix-a---geriatric-long-term-care-review-committee-(gltcrc)-report.pdf?sfvrsn=82eedab1_0; College of Occupational Therapists of Ontario, "2022 College Response to the Coroner's Report: Deaths from Power Recliner Lift Chairs," Feb. 2022, https://www.coto.org/docs/default-source/default-document-library/coto-coroners-report-response-deaths-from-power-recliner-lift-chairs.pdf?sfvrsn=f2d46c5d_2.

[5] Connors, Karol A., et al., "Adaptive Equipment Use by People with Motor Neuron Disease in Australia: A Prospective, Observational Consecutive Cohort Study," Disability and Rehabilitation: Assistive Technology, Jan. 2019, https://www.ncbi.nlm.nih.gov/pubmed/29082797.

[6] Itokazu, M., et al., "Analysis of Rising from a Chair after Total Knee Arthroplasty," Bulletin: Hospital for Joint Diseases, 1998, https://www.ncbi.nlm.nih.gov/pubmed/9725064.

[7] Pai, Yi-Chung, and Mark W. Rogers, "Speed Variation and Resultant Joint Torques During Sit-to-Stand," Archives of Physical Medicine and Rehabilitation, Oct. 1991, https://www.archives-pmr.org/article/0003-9993(91)90004-3/fulltext.

[8] Hughes, M. A., and M. L. Schenkman, "Chair Rise Strategy in the Functionally Impaired Elderly," Journal of Rehabilitation Research and Development, Oct. 1996, https://www.ncbi.nlm.nih.gov/pubmed/8895136; Kotake, Tomomitsu, et al., "An Analysis of Sit-to-Stand Movements," Archives of Physical Medicine and Rehabilitation, Oct. 1993, https://www.archives-pmr.org/article/0003-9993(93)90068-L/fulltext.

[9] Podsiadlo, D., and Richardson, S., "The Timed Up & Go: A Test of Basic Functional Mobility for Frail Elderly Persons," Journal of the American Geriatrics Society, 1991;39(2):142-148, doi:10.1111/j.1532-5415.1991.tb01616.x, https://pubmed.ncbi.nlm.nih.gov/1991946/.

PRIDE MOBILITY®

[10] Ng, S. S., and Hui-Chan, C. W., "The Timed Up & Go Test: Its Reliability and Association with Lower-Limb Impairments and Locomotor Capacities in People with Chronic Stroke," Archives of Physical Medicine and Rehabilitation, 2005;86(8):1641-1647, doi:10.1016/j.apmr.2005.01.011, https://pubmed.ncbi.nlm.nih.gov/16084820/.

[11] Morris, S., Morris, M. E., and Iansek, R., "Reliability of Measurements Obtained With the Timed Up & Go Test in People With Parkinson Disease," Physical Therapy, 2001;81(2):810-818, doi:10.1093/ptj/81.2.810, https://pubmed.ncbi.nlm.nih.gov/11175678/; Thompson, M., and Medley, A., "Performance of Individuals with Parkinson's Disease on the Timed Up & Go," Journal of Neurologic Physical Therapy, 1998;22(1):16-21, doi:10.1097/01253086-199822010-00009.

[12] Zampieri, C., Salarian, A., Carlson-Kuhta, P., Nutt, J., and Horak, F., "Assessing Mobility at Home in People with Early Parkinson's Disease Using an Instrumented Timed Up and Go Test," Parkinsonism & Related Disorders, 2011;17(4):277-280, doi:10.1016/j.parkreldis.2010.08.001, https://pubmed.ncbi.nlm.nih.gov/20801706/; Salarian, A., Horak, F., Zampieri, C., Carlson-Kuhta, P., Nutt, J., and Aminian, K., "iTUG, a Sensitive and Reliable Measure of Mobility," IEEE Transactions on Neural Systems and Rehabilitation Engineering, 2010;18(3):303-310, doi:10.1109/TNSRE.2010.2047606, https://pubmed.ncbi.nlm.nih.gov/20388604/.

[13] Shumway-Cook, A., Brauer, S., and Woollacott, M., "Predicting the Probability for Falls in Community-Dwelling Older Adults Using the Timed Up & Go Test," Physical Therapy, 2000;80(9):896-903, doi:10.1093/ptj/80.9.896, https://pubmed.ncbi.nlm.nih.gov/10960937/.

[14] Guralnik, J. M., et al., "Lower Extremity Function and Subsequent Disability: Consistency Across Studies, Predictive Models, and Value of Gait Speed Alone Compared with the Short Physical Performance Battery," The Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 2000;55(4):M221-M231, doi:10.1093/gerona/55.4.M221.

[15] Vasunilashorn, S., et al., "Use of the Short Physical Performance Battery Score to Predict Loss of Ability to Walk 400 Meters: Analysis From the InCHIANTI Study," The Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 2009;64A(2):223-229, doi:10.1093/gerona/gln022.

[16] Hoenig, H. M., and Rubenstein, L. Z., "Hospital-Associated Deconditioning and Dysfunction," Journal of the American Geriatrics Society, 1991;39(2):220-222, doi:10.1111/j.1532-5415.1991.tb01632.x.

[17] Chen, Y., Almirall-Sanchez, A., Mockler, D., Adrion, E., Dominguez-Vivero, C., and Romero-Ortuno, R., "Hospital-Associated Deconditioning: Not Only Physical, But Also Cognitive," International Journal of Geriatric Psychiatry, 2022;37(3), doi:10.1002/gps.5687, https://pubmed.ncbi.nlm.nih.gov/35142397/.

[18] Garber, S. L., "Wheelchair Cushions: A Historical Review," American Journal of Occupational Therapy, 1985;39(7):453-459, doi:10.5014/ajot.39.7.453; Eakin, P. A., Porter-Armstrong, A. P., and Stinson, M. D., "Pressure Mapping Systems: Reliability of Pressure Map Interpretation," Clinical Rehabilitation, 2003;17(5):504-511, doi:10.1191/0269215503cr643oa; Ferguson-Pell, M., and Cardi, M. D., "Prototype Development and Comparative Evaluation of Wheelchair Pressure Mapping System," Assistive Technology, 1993;5(2):78-91.

401 YORK AVENUE, DURYEA, PA 18642

570.655.5574 • TOLL FREE 800.800.8586

PRIDEMOBILITY.COM

Rev A/November 2026

INFINFB4980